

IN THE UNITED STATES

PATENT AND TRADEMARK OFFICE

APPLICANT: Sharat Singh *et al.*
APPLICATION NO.: 10/520,230
FILING DATE: February 02, 2005
TITLE: Methods and compositions for screening cell binding molecules
EXAMINER: Gary W. Counts
GROUP ART UNIT: 1641
ATTY. DKT. NO.: 114.00US/25237-12870

MAIL STOP AMENDMENT
COMMISSIONER FOR PATENTS
P.O. BOX 1450
ALEXANDRIA, VA 22313-1450

RESPONSE TO RESTRICTION REQUIREMENT

SIR:

Responsive to the Office Action (Restriction Requirement) dated June 29, 2007 received in the above-identified patent application, please consider the accompanying remarks. The Response is filed within one month of the mailing of the Office Action, therefore, no fees are believed due. However, the Commissioner is hereby authorized to charge to Deposit Account No. 19-2555 any other fees under 37 C.F.R. §§ 1.16, 1.17 and 1.21 which may be required by this paper, with the exception of the payment of the Issue Fee. Reconsideration of the application in view of the following remarks is respectfully requested.

REMARKS

STATUS OF THE CLAIMS

1. (Original) A method of determining the binding specificity of a ligand to a cell surface moiety, the method comprising the steps of:

(a) providing one or more cell types such that each cell type has a different membrane-anchored electrophoretic probe, each membrane-anchored electrophoretic probe having a membrane anchoring moiety connected by a cleavable linkage to an electrophoretic tag, said tag having distinct optical or electrophoretic properties with respect to electrophoretic tags of other cell types;

(b) combining the one or more cell types with a ligand, and

(c) exposing the cell types and ligand to conditions such that at least one cleavable linkage is cleaved in at least one membrane-anchored electrophoretic probe of cells having a cell surface moiety to which said ligand is bound, whereby one or more electrophoretic tags are released; and

(d) electrophoretically separating and identifying the one or more released electrophoretic tags, to determine the specificity of the ligand for the cell surface moieties of the cell types.

2. (Original) The method of claim 1, wherein said exposing comprises

attaching to each cell having a bound ligand, a proximity-dependent cleavage inducing group, such that each said probe is cleavable only by such a cleavage inducing group on the same cell surface as the probe, and

activating said cleavage inducing group.

3. (Original) The method of claim 2, wherein said probe is cleavable by a short-lived chemical species generated by a sensitizer group, and said exposing includes

attaching such a sensitizer group to each cell having a bound ligand, and

activating the sensitizer group.

4. (Original) The method of claim 3, wherein the sensitizer group is a photosensitizer.
5. (Original) The method of claim 3, wherein the sensitizer group is attached to the ligand.
6. (Original) The method of claim 3, wherein the sensitizer group is attached to a secondary molecule which forms a complex with the ligand.
7. (Original) The method of claim 6, wherein the ligand is an antibody, the sensitizer is conjugated to a secondary antibody immunospecific against the antibody, and said exposing includes adding the secondary antibody and conjugated sensitizer to the cells and bound antibody.
8. (Original) The method of claim 1, wherein said step of combining comprises isolating cell types having at least one ligand bound to at least one cell surface moiety.
9. (Original) The method of claim 8, wherein said cleavable linkage is chemically cleavable, photochemically cleavable, or enzymatically cleavable.
10. (Original) A method of identifying a cell surface antigen specific to substantially only one of a plurality of cell types, the method comprising the steps of:
 - (a) providing a plurality of cell types with cell surface antigens such that each cell type has a different membrane-anchored electrophoretic probe, the membrane-anchored electrophoretic probe having a membrane anchoring moiety connected by a cleavable linkage to an electrophoretic tag, said tag having distinct optical or electrophoretic properties with respect to electrophoretic tags of other cell types of the plurality;
 - (b) combining the one or more cell types with a candidate antibody;

(c) exposing the cell types and antibody to conditions such that at least one cleavable linkage is cleaved in at least one membrane-anchored electrophoretic probe of cells having a cell surface antigen to which said antibody is bound, whereby one or more electrophoretic tags are released;

(d) electrophoretically separating and determining the relative quantities of the one or more released electrophoretic tags to determine whether the candidate antibody binds to a cell surface antigen present on substantially only one of the plurality of cell types; and

(e) repeating steps (b)-(d) until a cell surface antigen specific to substantially only one of the plurality of cell types is identified.

11. (Original) The method of claim 10, wherein said exposing comprises

attaching to each cell having a bound antibody, a proximity-dependent cleavage inducing group, such that each said probe is cleavable only by such a cleavage inducing group on the same cell surface as the probe, and

activating said cleavage inducing group.

12. (Original) The method of claim 11, wherein said probe is cleavable by a short-lived chemical species generated by a sensitizer group, and said exposing includes

attaching such a sensitizer group to each cell having a bound antibody, and

activating the sensitizer group.

13. (Original) The method of claim 12, wherein the sensitizer group is a photosensitizer.

14. (Original) The method of claim 13, wherein the sensitizer group is attached to the antibody.

15. (Original) The method of claim 13, wherein the sensitizer is conjugated to a secondary antibody immunospecific against the antibody, and said exposing includes adding the secondary antibody and conjugated sensitizer to the cells and bound antibody.

16. (Original) The method of claim 10, wherein said step of combining comprises isolating cell types having at least one antibody bound to at least one cell surface antigen.

17. (Original) The method of claim 16, wherein said cleavable linkage is chemically cleavable, photochemically cleavable, or enzymatically cleavable.

18. (Original) The method of claim 10, wherein the determining of step (d) comprises measuring the area of tag peaks in an electropherogram of said released tags, and identifying a cell surface antigen specific to substantially only one of the plurality of cell types comprises identifying one test tag peak in said electropherogram that is at least 90% of the sum of the areas of all the test tag peaks in the electropherogram.

19. (Original) The method of claim 18, wherein said one test eTag peak is at least 95% of the sum of the areas of all the test eTag peaks in the electropherogram.

20. (Original) The method of claim 18, wherein identifying a cell surface antigen specific to substantially only one of the plurality of cell types comprises identifying one test tag peak in said electropherogram having an area that is at least twice the area of the next largest test tag peak in the electropherogram.

21. (Original) The method of claim 20, wherein said one test tag peak is at least four times the area of the next largest test tag peak in the electropherogram.

22. (Original) The method of claim 10, further comprising identifying said candidate antibody which binds to said cell surface antigen.

23. (Original) A method of determining the binding affinity of a compound for a cell surface antigen, the method comprising the steps of:

(a) providing one or more test cell-antibody pairs, each such pair comprising (i) a test cell having a membrane-anchored electrophoretic probe, the membrane-anchored electrophoretic probe having a membrane anchoring moiety connected by a cleavable linkage to an electrophoretic tag, said tag having distinct optical or electrophoretic properties with respect to electrophoretic tags of other test cells of the plurality, and (ii) at least one antibody specific for an cell surface antigen of the test cell of such pair, such cell surface antigen being different from cell surface antigens on other test cells of the plurality;

(b) combining the compound with the test cell-antibody pairs under conditions that permit the binding of the antibodies and the compound to one or more cell surface antigens recognized by said compound or antibodies;

(c) exposing the test cell-antibody pairs to conditions such that at least one cleavable linkage is cleaved in at least one membrane-anchored electrophoretic probe of cells having a cell surface antigen to which said antibody is bound, whereby one or more electrophoretic tags are released;

(d) electrophoretically separating the released tags; and

(e) determining the relative quantities of each of the one or more released electrophoretic tags to determine the binding affinity of the compound for the cell surface antigens.

24. (Original) The method of claim 23, wherein said exposing comprises

attaching to each cell having a bound antibody, a proximity-dependent cleavage inducing group, such that each said probe is cleavable only by such a cleavage inducing group on the same cell surface as the probe, and
activating said cleavage inducing group.

25. (Original) The method of claim 24, wherein said probe is cleavable by a short-lived chemical species generated by a sensitizer group, and said exposing includes

attaching such a sensitizer group to each cell having a bound antibody, and
activating the sensitizer group.

26. (Original) The method of claim 25, wherein the sensitizer group is a photosensitizer.

27. (Original) The method of claim 26, wherein the sensitizer group is attached to the antibody.

28. (Original) The method of claim 26, wherein the sensitizer is conjugated to a secondary antibody immunospecific against the antibody, and said exposing includes adding the secondary antibody and conjugated sensitizer to the cells and bound antibody.

29. (Original) The method of claim 23, further comprising the step of comparing the relative quantities obtained in step (c) with those obtained when steps (a), (c) and (d) are carried out in the absence of step (b).

30. (Original) A method of determining the binding specificity of a compound for an internalizing cell surface receptor, the method comprising the steps of:

(a) providing a plurality of test cell-antibody pairs, each such pair comprising (i) a test cell having a membrane-anchored electrophoretic probe, the membrane-anchored electrophoretic probe having a membrane anchoring moiety connected by a cleavable linkage to an electrophoretic tag

having distinct optical or electrophoretic properties with respect to electrophoretic tags of other test cells of the plurality, and (ii) at least one antibody effective to bind to an internalizing cell surface receptor of the test cell of such pair;

(b) combining the compound with the plurality of test cells under conditions that permit the endocytosis of complexes that form between the compound and one or more of the internalizing cell surface receptors;

(c) combining with the test cells the plurality of antibodies under conditions that permit binding to internalizing cell surface receptors;

(d) exposing the test cell-antibody pairs to conditions such that at least one cleavable linkage is cleaved in at least one membrane-anchored electrophoretic probe of cells having an internalizing cell surface receptor to which said antibody is bound, whereby one or more electrophoretic tags are released;

(e) electrophoretically separating the released tags; and

(f) determining the relative quantities of each of the one or more released electrophoretic tags to determine the specificity of the compound for the plurality endocytosing cell surface receptors.

31. (Original) The method of claim 30, wherein said exposing comprises

attaching to each cell having a bound antibody, a proximity-dependent cleavage inducing group, such that each said probe is cleavable only by such a cleavage inducing group on the same cell surface as the probe, and

activating said cleavage inducing group.

32. (Original) The method of claim 30, wherein the internalized cell surface receptor of each test cell is different from internalizing cell surface receptors on other test cells of the plurality.

33. (Original) The method of claim 30, wherein said probe is cleavable by a short-lived chemical species generated by a sensitizer group, and said exposing includes

attaching such a sensitizer group to each cell having a bound antibody, and
activating the sensitizer group.

34. (Original) The method of claim 33, wherein the sensitizer group is a photosensitizer.

35. (Original) The method of claim 33, wherein the sensitizer group is attached to the antibody.

36. (Original) The method of claim 33, wherein the sensitizer is conjugated to a secondary antibody immunospecific against the antibody, and said exposing includes adding the secondary antibody and conjugated sensitizer to the cells and bound antibody.

37. (Original) The method of claim 30, further comprising the step of comparing the relative quantities obtained in step (f) with those obtained when steps (a) and (c)-(e) are carried out in the absence of step (b).

Response to Restriction Requirement

In the Office Action, the Examiner restricted the claims to 4 groups. The Examiner therein required election of one of the following groups of claims:

Group 1, claims 1-9, drawn to a method determining the binding specificity of a ligand to a cell surface moiety;

Group 2, claims 10-22, drawn to a method of identifying a cell surface antigen specific to substantially only one of a plurality of cell types;

Group 3, claims 23-29, drawn to a method of determining the binding affinity of a compound for a cell surface antigen; and

Group 4, claims 30-37, drawn to a method of determining the binding specificity of a compound for an internalizing cell surface receptor.

Applicants provisionally elect Group 2 comprising claims 10-22 without traverse. Applicants reserve the right to pursue the unselected subject matter in one or more related divisional, continuation, or continuation-in-part applications.

CONCLUSION

Consideration of the claims is respectfully requested, and a notice of allowance is earnestly solicited. If the Examiner has any questions concerning this Amendment, the Examiner is invited to telephone Applicants' representative at (650) 335-7818.

Respectfully submitted,
Sharat Singh *et al.*

Dated: July 25, 2007

By: /Narinder Banait/
Narinder S. Banait, Reg. No. 43,482
Fenwick & West LLP
801 California Street
Mountain View, CA 94041
Telephone: 650-335-7818
Facsimile: 650-938-5200
E-mail: nbanait@fenwick.com